

TRIVIAL AND SIMPLE SPECTRUM FOR $SL(2, \mathbb{R})$ COCYCLES WITH FREE BASE AND FIBER DYNAMICS

MÁRIO BESSA AND PAULO VARANDAS

ABSTRACT. We prove that there exists a residual subset $\mathcal{R} \subset \text{Aut}_\mu(M) \times L^\infty(M, SL(2, \mathbb{R}))$, with respect to the product topology, i.e., weak topology $\times L^\infty$, such that any element (f, A) in $\mathcal{R}$ has trivial spectrum or it is hyperbolic. Then, we prove prevalence of trivial spectrum near the dynamical cocycle of an area-preserving map and also for generic cocycles in $\text{Aut}_\mu(M) \times L^p(M, SL(2, \mathbb{R}))$. On the other hand, if $\mathcal{A} \subset \text{Diff}^1(M)$ denotes the set of C^1 -Anosov diffeomorphisms and $f \in \mathcal{A}$ there exists $\nu > 0$ and an open neighborhood V of f such that open and densely in dominated cocycles in $V \times C^\nu(M, SL(2, \mathbb{R}))$ the pair (f, A) has simple spectrum with respect to the unique maximal entropy measure μ_f .

1. INTRODUCTION

Let M denote a compact d -dimensional manifold ($d \geq 2$) endowed with a distance $d(\cdot, \cdot)$, μ the volume-measure associated to the volume form on M , $\text{Aut}_\mu(M)$ the space of the automorphisms of M (i.e. bi-measurable bijections preserving μ) and $\text{Homeo}_\mu(M)$ the space of homeomorphism $f: M \rightarrow M$ which leave invariant the measure μ . We endow $\text{Aut}_\mu(M)$ with the *weak topology* denoted by $\mathcal{W}$ (i.e. $f_n \rightarrow f$ if and only if $\mu(f_n(K) \triangle f(K)) \rightarrow 0$ for all measurable sets $K \subset M$) and endow $\text{Homeo}_\mu(M)$ with the C^0 topology. Let $A: M \rightarrow SL(2, \mathbb{R})$ be a continuous map when we consider the distance d in M and the uniform operator norm on the $SL(2, \mathbb{R})$. Let $r \in \mathbb{N}_0$ and $\nu \geq 0$ be such that $r + \nu > 0$ and let $C^{r+\nu}(M, SL(2, \mathbb{R}))$ denote the space of C^r cocycles $A: M \rightarrow SL(2, \mathbb{R})$ such that $D^r A$ is a C^ν -Hölder continuous. Let $L^\infty(M, SL(2, \mathbb{R}))$ denote the space of all essentially bounded maps $A: M \rightarrow SL(2, \mathbb{R})$ endowed with the L^∞ -norm defined by $\|A - B\|_\infty = \text{esssup} \|A(x) - B(x)\|$. Given $A \in L^\infty(M, SL(2, \mathbb{R}))$ we will denote, by a slight abuse of notation, by *cocycle* the skew-product

$$F_A: \begin{array}{ccc} M \times \mathbb{R}^2 & \longrightarrow & M \times \mathbb{R}^2 \\ (x, v) & \mapsto & (f(x), A(x) \cdot v), \end{array}$$

whose joint base and fiber dynamics is given by $F_A^n(x, v) = (f^n(x), A^n(x) \cdot v)$ where $A^n(x) = A(f^{n-1}(x)) \circ \dots \circ A(f(x)) \circ A(x)$. If μ is an f -invariant probability measure such that $\log \|A^\pm\| \in L^1(\mu)$ then it follows from Oseledets theorem (see e.g. [2]) that for μ -almost every x there exists the *largest Lyapunov exponent* defined by the limit

$$\lambda^+(f, A, x) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log \|A^n(x)\|,$$

satisfying $\lambda^+(f, A, f(x)) = \lambda^+(f, A, x)$ and it is a non-negative measurable function of x . Moreover, for μ -a.e. point $x \in M$ with $\lambda(f, A, x) \neq 0$, there is a A -invariant splitting of the bundle over x , $E_x^u \oplus E_x^s$ which varies measurably with x and such that, if $u \in E_x^u \setminus \{\tilde{0}\}$ and $s \in E_x^s \setminus \{\tilde{0}\}$, then

$$\lim_{n \rightarrow \pm\infty} \frac{1}{n} \log \|A^n(x) \cdot u\| = \lambda^+(f, A, x) \quad \text{and} \quad \lim_{n \rightarrow \pm\infty} \frac{1}{n} \log \|A^n(x) \cdot s\| = \lambda^-(f, A, x) = -\lambda^+(f, A, x).$$

2. TRIVIAL AND SIMPLE SPECTRUM

In the recent years several results about the generic behavior of $SL(2, \mathbb{R})$ cocycles have been proved. We refer to result of Cong [7] where it is proved the L^∞ genericity of hyperbolic behavior among bounded cocycles when we keep the base dynamics and perturb the fiber dynamics in the L^∞ -sense. Moreover, Bochi and Fayad [4] proved that the trivial Lyapunov spectrum prevails in a C^0 -generic way when we keep the fiber dynamics and allows perturbation in the base dynamics.

In this note we intend to discuss the problem of knowing what is the prevalent behaviour when we consider two degrees of freedom, i.e., perturbation of both base and fiber dynamics. In what follows, we endow the product

space $\text{Aut}_\mu(M) \times L^\infty(M, SL(2, \mathbb{R}))$ with the product topology $\mathcal{W} \times L^\infty$. Moreover, we let (M, μ) be a non-atomic Lebesgue space. The following result is very simple consequence of [4, Proposition 1.7] which we present in order to contextualize with the object of our study.

Theorem 2.1. ([4, Proposition 1.7]) *There exists a residual subset $\mathcal{R} \subset \text{Aut}_\mu(M) \times L^\infty(M, SL(2, \mathbb{R}))$ such that any element $(f, A) \in \mathcal{R}$ has trivial spectrum or else the cocycle (f, A) is uniformly hyperbolic.*

Proof. We consider the function

$$\begin{aligned} \Lambda: \quad \text{Aut}_\mu(M) \times L^\infty(M, SL(2, \mathbb{R})) &\longrightarrow [0, +\infty[\\ (f, A) &\mapsto \int_M \lambda^+(f, A, x) d\mu(x) \end{aligned}$$

Take the residual $\mathcal{R}_1 \subset L^\infty(M, SL(2, \mathbb{R}))$ given in [4, Proposition 1.7]. Then, for every $A \in \mathcal{R}_1$ we have that (f, A) is uniformly hyperbolic for any $f \in \text{Aut}_\mu(M)$ or else there exists a residual $\mathcal{R}_2$ (w.r.t. $\mathcal{W}$) such that any $f \in \mathcal{R}_2$ satisfies $\Lambda(f, A) = 0$. Define $\mathcal{R} = \mathcal{R}_1 \times \mathcal{R}_2$. $\square$

Remark 2.2. If we assume that M is a compact Hausdorff space and μ is a Borel regular measure, then previous result also holds for $\text{Aut}_\mu(M) \times C^0(M, SL(2, \mathbb{R}))$ instead of $\text{Aut}_\mu(M) \times L^\infty(M, SL(2, \mathbb{R}))$ and the C^0 topology instead of the L^∞ one.

3. TRIVIAL SPECTRUM

Here, we begin by proving that the trivial spectrum is prevalent for the dynamical cocycle of an area-preserving diffeomorphism near any element of $\text{Homeo}_\mu(M)$. Let $\text{Diff}_\mu^r(M)$ stands for the set of area-preserving diffeomorphisms on surfaces.

Theorem A. *For a C^0 -dense subset $\mathcal{D} \subset \text{Homeo}_\mu(M)$ of C^2 diffeomorphisms and $\varepsilon > 0$ we have that, for $f \in \mathcal{D}$, there is a C^0 -neighborhood $\mathcal{U}_f$ of f and a residual subset $\mathcal{R} \subset \mathcal{U}_f$ such that $\int_M \lambda^+(h, Df, x) d\mu(x) < \varepsilon$, for all $h \in \mathcal{R}$.*

Proof. Using [11] we know that C^1 -close to any element in $\text{Diff}_\mu^1(M)$ there is an Anosov map or else one map that exhibits dense elliptic periodic orbits. By ([13]) $\text{Diff}_\mu^r(M)$ ($r \geq 2$) is C^1 -dense in $\text{Diff}_\mu^1(M)$ and by ([10]) $\text{Diff}_\mu^1(M)$ is C^0 -dense in $\text{Homeo}_\mu(M)$. By performing an arbitrarily small C^0 -perturbation of an Anosov in $\text{Diff}_\mu^1(M)$ we can obtain an element in $\text{Diff}_\mu^1(M)$ C^1 -far from the C^1 -open subset of Anosovs, thus having an element in $\text{Diff}_\mu^1(M)$ exhibiting dense elliptic periodic orbits. Therefore, we obtain a C^0 -dense subset $\mathcal{D}$ of $\text{Homeo}_\mu(M)$ with dense elliptic periodic orbits and of class C^2 . Moreover, $\mathcal{D}$ can be chosen such that at least one of its (dense) periodic orbits has non-zero second derivative at some point of its orbit.

In conclusion, we have $f \in \mathcal{D}$ with the dynamical cocycle $Df: M \rightarrow SL(2, \mathbb{R})$ of class C^1 and, moreover, f has a persistent¹ elliptic periodic point $p = f^n(p)$ such that $D(Df_{f^i(p)})$ is non-zero, for some $i \in \{0, 1, \dots, n-1\}$. We are in the conditions of [4, Corollary 5] (see also [4, Theorem 4]) and so we obtain a neighborhood $\mathcal{U}_f \subset \text{Homeo}_\mu(M)$ of f and a residual subset $\mathcal{R} \subset \mathcal{U}_f$ such that $\int_M \lambda^+(h, Df, x) d\mu(x) < \varepsilon$ for all $h \in \mathcal{R}$ and the theorem is proved. $\square$

Now, we also obtain prevalence of trivial spectrum if we consider the L^p -measurable $SL(2, \mathbb{R})$ cocycles endowed with the L^p -norm (c.f. [1]). The next result is a direct consequence of Arbieto & Bochi theorem [1].

Corollary 3.1. *There exists a residual subset $\mathcal{R} \subset \text{Aut}_\mu(M) \times L^p(M, SL(2, \mathbb{R}))$ such that any element $(f, A) \in \mathcal{R}$ has trivial spectrum.*

Proof. We consider the function endowed with the product topology, $\mathcal{W} \times L^p$ -norm, in its domain:

$$\begin{aligned} \Lambda: \quad \text{Aut}_\mu(M) \times L^p(M, SL(2, \mathbb{R})) &\longrightarrow [0, +\infty[\\ (f, A) &\mapsto \int_M \lambda^+(f, A, x) d\mu(x) \end{aligned}$$

By [4, Proposition A.2] we know that, for any fixed $A \in L^p(M, SL(2, \mathbb{R}))$, we have that the integrated Lyapunov exponent function $\Lambda(\cdot, A)$ is upper semicontinuous w.r.t. $\mathcal{W}$. Moreover, by [1, Theorem 2], for any fixed $f \in \text{Aut}_\mu(M)$ we have that $\Lambda(f, \cdot)$ is upper semicontinuous with respect to the L^p norm. Hence, since we consider the product topology it is easy to show that Λ is an upper semicontinuous map. Thus, for every $\varepsilon > 0$ there exists a

¹A periodic point is *persistent* for $h \in \text{Homeo}_\mu(M)$ if for every $\varepsilon > 0$ there is $\delta > 0$ such that if $\hat{h}$ is an homeomorphism δ - C^0 -close to h then $\hat{h}$ has a periodic point $\hat{p}$ of period n which is ε -close to p . Since our elliptic point is an isolated fixed point, for the return map, of Poincaré-Lefschetz index different from 1 it is necessarily persistent.

neighborhood U of (f_0, A_0) such that $\Lambda(f, A) \leq \Lambda(f_0, A_0) + \varepsilon$ for all $(f, A) \in U$. Now we claim that if (f_0, A_0) is a continuity point of Λ then $\Lambda(f_0, A_0) = 0$. Assume, by contradiction, that $\Lambda(f_0, A_0) > 0$. Then, there exists a small $\varepsilon > 0$ so that $\Lambda(f, A) \geq \Lambda(f_0, A_0)/2 > 0$ for all $(f, A) \in \text{Aut}_\mu(M) \times L^p(M, SL(2, \mathbb{R}))$ such that f is ε -close to f_0 (w.r.t. the topology $\mathcal{W}$) and $\|A - A_0\|_{L^p} < \varepsilon$. However, we can perturb f_0 obtaining an ergodic f_1 (cf. [8]), and, since we are in the conditions of [4, Theorem 1], there exists $\tilde{A}_1 \in L^p(M, SL(2, \mathbb{R}))$ that is ε - L^p -close to A_0 and such that $\Lambda(\tilde{f}_1, \tilde{A}_1) = 0$ which leads to a contradiction. This proves our claim. Finally, since the set of continuity points of a semicontinuous function is a residual set the corollary is now proved. $\square$

4. SIMPLE SPECTRUM

From now on we will consider more regular cocycles with a fiber-bunching assumption called *domination* as we now describe. Denote by $\mathcal{A} \subset \text{Diff}^1(M)$ the space of C^1 -Anosov diffeomorphisms on M . Given $f \in \mathcal{A}$ there are constants $C = C_f > 0$ and $\theta = \theta_f \in (0, 1)$, and a Df -invariant splitting $TM = E_f^s \oplus E_f^u$ such that $\|Df_{E^s}^n\| \leq C\theta^n$ and $\|(Df_{E^u}^n)^{-1}\| \leq C\theta^n$ for all $n \geq 1$. In fact it is well known that $\mathcal{A}$ is an open set and that the constants $C > 0$ and $\theta_f \in (0, 1)$ can be chosen in a way that for all $g \in \mathcal{A}$ sufficiently close to f there exists a Dg -invariant splitting $TM = E_g^s \oplus E_g^u$, $\|Dg_{E^s}^n\| \leq C\theta^n$ and $\|(Dg_{E^u}^n)^{-1}\| \leq C\theta^n$ for all $n \geq 1$. Given a constant $\nu > 0$, we say that a ν -Hölder continuous cocycle A is *dominated* for f if it satisfies $\|A(x)\| \|A(x)^{-1}\| \theta^\nu < 1$ for all $x \in M$. This is clearly an open condition. Now, since Anosov are structurally stable for any $f \in \mathcal{A}$ there exists $\gamma \in (0, 1)$ and a γ -Hölder continuous homeomorphism h_g close to the identity and such that $g \circ h_g = h_g \circ f$. Let $\eta : \mathcal{A} \rightarrow (0, 1]$ be a continuous function such that any $f \in \mathcal{A}$ is $\eta(f)$ -Hölder conjugate to all sufficiently close maps (see e.g. [9] for details on regularity of conjugacies) and let $AC_D^\eta(M, SL(2, \mathbb{R})) \subset \mathcal{A} \times C^0(M, SL(2, \mathbb{R}))$ to be the open set of pairs (f, A) such that f is an Anosov diffeomorphism, A is a $\eta(f)$ -Hölder continuous cocycle and it is dominated for f .

It is well known that these Anosov diffeomorphisms are ergodic, hence transitive. In particular for every $f \in \mathcal{A}$ there exists a unique maximal entropy measure μ_f , and it has local product structure as we now describe. Recall that local stable and local unstable manifolds are C^1 -embedded submanifolds of M with the property that $W_{\text{loc}}^s(x)$ and $W_{\text{loc}}^u(x)$ vary continuously with x and there exists $\delta > 0$ is small such that for any $x \in M$ and $y, z \in B(x, \delta)$ the intersection $[y, z] := W_{\text{loc}}^u(y) \cap W_{\text{loc}}^s(z) \neq \emptyset$ consists of a unique point. Hence, there exists $N_x^u(\delta) \subset W_{\text{loc}}^u(x)$ a u -neighborhood of x and $N_x^s(\delta) \subset W_{\text{loc}}^s(x)$ an s -neighborhood of x and a neighborhood $N_\delta(x)$ of x in M such that the map $\Upsilon_x : N_\delta(x) \rightarrow N_x^u(\delta) \times N_x^s(\delta)$ given by $\Upsilon_x(y) = ([x, y], [y, x])$ is a homeomorphism. An f -invariant probability measure μ has *local product structure* if for any $x \in \text{supp}(\mu)$ ($\text{supp}(\mu)$ stands for the support of the measure μ) and a small $\delta > 0$ the measure $\mu|_{N_x(\delta)}$ is equivalent to the product measure $\mu_x^u \times \mu_x^s$, where μ_x^i denotes the conditional measure of $(\Psi_x)_*(\mu|_{N_x(\delta)})$ on $N_x^i(\delta)$, for $i \in u, s$. See e.g. [6] for details. We study Lyapunov exponent with respect to the maximal entropy measure:

$$\begin{aligned} \Lambda : \quad AC_D^\eta(M, SL(2, \mathbb{R})) &\longrightarrow \mathbb{R} \\ (f, A) &\longmapsto \int_M \lambda^+(f, A, x) d\mu_f \end{aligned}$$

Let $f : M \rightarrow M$ and $g : N \rightarrow N$ be invertible measurable maps and measurable h -conjugated, say $g \circ h = h \circ f$, for an invertible measurable map $h : M \rightarrow N$. The cocycle A over f and the cocycle B over g are *equivalent* if there exists a measurable tempered map $L : M \rightarrow SL(2, \mathbb{R})$ such that the cohomology equations holds: $A(h^{-1}(x)) = L(g(x))^{-1} B(x) L(x)$ for $x \in N$ (c.f. Chapter 4 in [2]). We are now in a position to state our last main result.

Theorem B. *There exists an open and dense subset $\mathcal{R} \subset AC_D^\eta(M, SL(2, \mathbb{R}))$ such that any element $(f, A) \in \mathcal{R}$ has simple spectrum with respect to the maximal entropy measure μ_f .*

Proof. Let A be a dominated cocycle for f such that (f, A) has simple spectrum, that is, $A \in \mathcal{S}_f$. We prove that there exists a small neighborhood $\mathcal{U}$ of f such that for all $g \in \mathcal{U}$ it holds that (g, A) has simple spectrum. Fix $\varepsilon > 0$ arbitrary small and $\|f - g\| < \varepsilon$. Then there exists a $\eta(f)$ -Hölder continuous homeomorphism h_g that is close to the identity and such that $g \circ h_g = h_g \circ f$. Since μ has local product structure for all Anosov diffeomorphisms in $\mathcal{A}$ it follows from [5, 12] that for any fixed $f \in \mathcal{A}$ there exists an open and dense subset $\mathcal{S}_f \subset C^{\eta(f)}(M, SL(2, \mathbb{R}))$ such that, for all $A \in \mathcal{S}_f$, the corresponding cocycle has simple spectrum for μ_f -almost every x . So, there exists a $C^{\eta(f)}$ -open neighborhood $\mathcal{V} \subset \mathcal{S}_f$ of A such that for all $\tilde{A} \in \mathcal{V}$ it holds $\|A - \tilde{A}\|_\nu \ll \varepsilon$ and $(f, \tilde{A})$ has simple spectrum with respect to μ_f . Observe also that, since domination is an open condition, if a cocycle A is dominated for f then it is dominated for all $g \in \mathcal{A}$ sufficiently close to f . Now, we claim that the map

$$g \mapsto \tilde{A} = A \circ h_g^{-1} \in \mathcal{V}$$

is well defined, continuous and surjective provided that $\mathcal{V}$ is small. Since $h_g^{-1} \in C^{\eta(f)}$ then it follows that $\tilde{A}$ is a $C^{\eta(f)}$ -Hölder continuous cocycle. Moreover, $\|\tilde{A} - A\| = \sup_x \|A(x) - A(h_g^{-1}(x))\| < \varepsilon$, since the second term can be taken arbitrary small provided that g is sufficiently close to f , and also $\|\tilde{A} - A\|_{\eta}$ can be taken arbitrary small using an easy argument. Using the continuity of the conjugacy it follows that the previous map is continuous and well defined as claimed. The surjection is immediate.

Now, since there is a unique maximal entropy measure for both f and g then it is preserved by topological conjugacy and so $(h_g^{-1})_*\mu_g = \mu_f$. Thence, the preimage is an open subset and thus (g, A) has simple spectrum for μ_g -almost every x . We have proven that, for any $A \in C^{\eta(f)}(M, SL(2, \mathbb{R}))$, which has simple spectrum for some (f, μ_f) the set of g 's such that (g, A) has simple spectrum for μ_g -almost every x is open. The statement of the theorem follows directly. $\square$

Remark 4.1. The previous result also extends from Anosov to transitive Axiom A diffeomorphisms satisfying the transversality condition, since these are structurally stable and locally bi-Hölder conjugate and $SL(d, \mathbb{R})$ cocycles for $d \geq 2$.

ACKNOWLEDGEMENTS

MB was partially supported by National Funds through FCT - “Fundação para a Ciência e a Tecnologia”, project PEst-OE/MAT/UI0212/2011 and also the project PTDC/MAT/099493/2008. PV was partially supported by CNPq and FAPESB. The authors would like to thanks Jairo Bochi for suggestions that improved the presentation of the paper.

REFERENCES

- [1] A. Arbieto and J. Bochi, L^p -generic cocycles have one-point Lyapunov spectrum, *Stochastics and Dynamics*, 3 (2003), 73–81. Corrigendum. *ibid*, 3 (2003), 419–420.
- [2] L. Barreira and Ya. Pesin, Nonuniform Hyperbolicity, *Encyclopedia of Mathematics and Its Applications 115*, Cambridge University Press, 2007.
- [3] J. Bochi. Genericity of zero Lyapunov exponents. *Erg. Theor. Dyn. Syst.*, 22:1667–1696, 2002.
- [4] J. Bochi and B. Fayad. Dichotomies between uniform hyperbolicity and zero Lyapunov exponents for $SL(2, \mathbb{R})$ cocycles. *Bull. Braz. Math. Soc.*, 37:307–349, 2006.
- [5] C. Bonatti and M. Viana. Lyapunov exponents with multiplicity 1 for deterministic products of matrices. *Ergod. Th. & Dynam. Sys.*, 24:1295–1330, 2004.
- [6] R. Bowen. Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms, *Lecture Notes in Mathematics*, 470, Springer, 2008.
- [7] N. Cong. A generic bounded linear cocycle has simple Lyapunov spectrum. *Erg. Theor. Dyn. Syst.*, 25:1775–1797, 2005.
- [8] P.R. Halmos. Lectures on Ergodic Theory. The Mathematical Society of Japan, 1956.
- [9] A. Katok and B. Hasselblat. Introduction to the Modern Theory of Dynamical Systems. *Encyclopedia of Mathematics and its Applications*, 1996.
- [10] Yong-Geun Oh. C^0 -coerciveness of Moser’s problem and smoothing area preserving homeomorphisms, Preprint ArXiv. math.DS/0601183, 2006.
- [11] S. Newhouse. Quasi-elliptic periodic points in conservative dynamical systems. *Am. J. Math.*, 99:1061–1087, 1977.
- [12] M. Viana, Almost all cocycles over any hyperbolic system have nonvanishing Lyapunov exponents. *Ann. of Math.* (2) 167 (2008), no. 2, 643–680.
- [13] E. Zehnder, Note on smoothing symplectic and volume-preserving diffeomorphisms, In *Proc. III Latin Amer. School of Math., Inst. Mat. Pura Aplicada CNPq, Rio de Janeiro, 1976*, volume 597 *Lecture Notes in Math.*, 828–854. Springer, Berlin, 1977.

MÁRIO BESSA, DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DA BEIRA INTERIOR, RUA MARQUÊS D’ÁVILA E BOLAMA, 6201-001 COVILHÃ, PORTUGAL.

E-mail address: `bessa@fc.up.pt`

PAULO VARANDAS, DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA BAHIA, AV. ADEMAR DE BARROS S/N, 40170-110 SALVADOR, BRAZIL.

E-mail address: `paulo.varandas@ufba.br`